

AUSTRALIAN PRODUCT INFORMATION – PONSTAN[®] CAPSULES (MEFENAMIC ACID)

1. NAME OF THE MEDICINE

Mefenamic acid

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

PONSTAN capsules contain the active ingredient mefenamic acid 250 mg/capsule.

Excipient(s) with known effect

- Lactose monohydrate

For the full list of excipients, see Section 6.1 List of excipients.

3. PHARMACEUTICAL FORM

Hard capsule.

PONSTAN capsules have an opaque aqua blue cap and opaque ivory body with “PARKE DAVIS” printed in black on both the body and cap.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment of primary dysmenorrhoea and primary menorrhagia.

Short-term relief of mild to moderate pain such as dental pain and soft tissue pain.

4.2 Dose and method of administration

Undesirable effects may be minimised by using the lowest effective dose for the shortest duration necessary to control symptoms.

Patients on long term treatment should be reviewed regularly with regards to efficacy, risk factors and ongoing need for treatment.

Dysmenorrhoea

2 capsules (500 mg) three times daily with meals from the onset of pain and continued for the usual duration of pain.

Menorrhagia

2 capsules (500 mg) three times daily with meals and from the onset of menses and continued according to the judgement of the physician. Therapy should not be continued for more than 7 days except on the advice of a physician.

Other Indications

2 capsules (500 mg) three times daily, with meals.

4.3 Contraindications

1. Patients showing evidence of chronic inflammation and/or active ulceration of either the upper or lower gastrointestinal tract and patients with pre-existing renal disease.
2. Patients in whom aspirin and/or other NSAIDs have induced symptoms of bronchospasm, allergic rhinitis or urticaria, because the potential exists for cross-sensitivity.
3. Patients with impaired renal function.
4. Patients with severe hepatic impairment.
5. Patients previously experiencing diarrhoea on taking this drug.
6. Patients who have previously exhibited hypersensitivity to mefenamic acid or any of the components of PONSTAN (See section 6.1 - List of excipients).
7. Patients with severe heart failure.
8. Treatment of peri-operative pain in the setting of coronary artery bypass graft (CABG) surgery.
9. Children under 14 years of age.

4.4 Special warnings and precautions for use

The use of PONSTAN with concomitant systemic non-aspirin NSAIDs, including cyclooxygenase-2 (COX-2) inhibitors, should be avoided. Concomitant use with one or more systemic NSAIDs may increase frequency of gastrointestinal ulcers and bleeding.

Cardiovascular Effects

Cardiovascular Thrombotic Events: NSAIDs may cause an increased risk of serious cardiovascular (CV) thrombotic events, myocardial infarction and stroke, which can be fatal. This risk may increase with dose or duration of use. The relative increase of this risk appears to be similar in those with or without known CV disease or CV risk factors. However, patients with CV disease, history of atherosclerotic CV disease or CV risk factors may also be at greater risk in terms of absolute incidence, due to their increased rate at baseline. To minimise the potential risk for an adverse CV event in patients treated with mefenamic acid, especially in patients with CV risk factors, the lowest effective dose should be used for the shortest duration possible. Physicians and patients should remain alert for the development of such events, even in the absence of previous CV symptoms. Patients should be informed

about the signs and/or symptoms of serious CV toxicity and the steps to take if they occur (See section 4.3 - Contraindications).

There is no consistent evidence that concurrent use of aspirin mitigates the increased risk of serious CV thrombotic events associated with NSAID use.

Hypertension: NSAIDs may lead to the onset of new hypertension or worsening of pre-existing hypertension and patients taking anti-hypertensives with NSAIDs may have an impaired anti-hypertensive response. Caution is advised when prescribing NSAIDs to patients with hypertension. Blood pressure should be monitored closely during initiation of NSAID treatment and at regular intervals throughout the course of therapy.

Heart Failure: Fluid retention and oedema have been observed in some patients taking NSAIDs, including mefenamic acid. Therefore, PONSTAN should be used with caution in patients with compromised cardiac function and other conditions predisposing to, or worsened by, fluid retention. Patients with pre-existing heart failure or hypertension should be closely monitored.

Gastrointestinal Effects

NSAIDs, including mefenamic acid, can cause serious, potentially fatal gastrointestinal (GI) adverse events including inflammation, bleeding, ulceration and perforation of the stomach, small intestine or large intestine. The frequency of such events may increase with dose or duration of use, but can occur at any time without warning.

Upper GI ulcers, gross bleeding or perforation caused by NSAIDs occur in approximately 1% of patients treated for 3-6 months and in about 2-4% of patients treated for one year. These trends continue with longer duration of use, increasing the likelihood of developing a serious GI event at some time during the course of therapy. However, even short term therapy is not without risk.

Patients most at risk of developing GI complications with NSAIDs are the elderly; patients with CV disease; patients using concomitant, corticosteroids, antiplatelet drugs (such as aspirin), or selective serotonin reuptake inhibitors (SSRIs); patients with a history of, or active, GI disease (such as ulceration, bleeding or inflammatory conditions); and patients ingesting alcohol or with a history of smoking and alcoholism. Mefenamic acid should be used with caution in these patients (See section 4.3 - Contraindications). When GI bleeding or ulceration occurs in patients receiving PONSTAN, the drug should be withdrawn immediately. Physicians should inform patients about the signs and/or symptoms of serious GI toxicity and what steps to take if they occur.

If diarrhoea occurs, PONSTAN should be promptly discontinued. Symptoms may recur in certain patients following subsequent exposure.

Severe Skin Reactions

NSAIDs may very rarely cause serious cutaneous adverse events such as exfoliative dermatitis, toxic epidermal necrolysis (TEN) and Stevens-Johnson syndrome (SJS), which can be fatal and occur without warning. These serious adverse events are idiosyncratic and are independent of dose or duration of use. Patients should be advised of the signs and symptoms of serious skin reactions and to consult their doctor at the first appearance of a skin

rash, mucosal lesion or any other sign of hypersensitivity. If this occurs, the drug should be promptly discontinued.

PONSTAN may cause an exacerbation of chronic urticaria in patients with this disease.

Haematologic Effects

PONSTAN 500 mg and aspirin 650 mg, four times a day, both caused significant further lowering of the prothrombin concentration (PONSTAN 3.48% and aspirin 2.75%) in patients in whom the concentration has been initially lowered by anticoagulant therapy. Caution should therefore be exercised in administering PONSTAN to patients on anticoagulant therapy, such as warfarin, and should not be given when prothrombin concentration is in the range of 10% to 20% of normal. Careful monitoring of blood coagulation factors is recommended (See section 4.5 - Interactions with other medicines and other forms of interactions).

Use with Oral Anticoagulants

The concomitant use of NSAIDs, including PONSTAN, with oral anticoagulants increases the risk of GI and non-GI bleeding and should be given with caution. Oral anticoagulants include warfarin/coumarin-type and novel oral anticoagulants (e.g. apixaban, dabigatran, rivaroxaban). Anticoagulation/INR should be monitored in patients taking a warfarin/coumarin-type anticoagulant (See section 4.5 - Interactions with other medicines and other forms of interactions).

Other Effects

PONSTAN should be used with caution in known asthmatics.

As no data presently exist concerning the effect of PONSTAN, if any, on the efficacy of intrauterine contraceptive devices, physicians should be alert to the possibility of a reduction in contraceptive efficacy in women with an IUCD taking PONSTAN.

Use in Hepatic Impairment

PONSTAN should be used with caution in patients with hepatic impairment.

As with other NSAIDs, borderline elevations of liver function tests may occur in up to 15% of patients. These elevations may progress, may remain essentially unchanged, or may be transient with continued therapy. Meaningful (3 times the upper limit of normal) elevations of SGPT or SGOT occurred in controlled clinical trials in less than 1% of patients.

Physicians and patients should remain alert for hepatotoxicity. Patients should be informed about the signs and/or symptoms of hepatotoxicity. A patient with symptoms and/or signs suggesting liver dysfunction (e.g. nausea, fatigue, lethargy, pruritis, jaundice, abdominal tenderness in the right upper quadrant and “flu-like” symptoms), or in whom an abnormal liver test has occurred, should be evaluated for evidence of the development of a more severe hepatic reaction while on therapy with mefenamic acid.

Severe hepatic reactions, including jaundice and cases of fatal hepatitis, have been reported with other NSAIDs. Although such reactions are rare, if abnormal liver tests persist or worsen, if clinical signs and symptoms consistent with liver disease develop, or if systemic manifestations occur (e.g. eosinophilia, rash, etc.), PONSTAN should be discontinued.

Use in Renal Impairment

As with other NSAIDs, long-term administration of PONSTAN to animals has resulted in renal papillary necrosis and other abnormal renal pathology. In humans, there have been reports of acute interstitial nephritis with haematuria, proteinuria, glomerulitis, papillary necrosis and occasionally nephrotic syndrome.

A second form of renal toxicity has been seen in patients with pre-renal conditions leading to a reduction in renal blood flow or blood volume, where the renal prostaglandins have a supportive role in the maintenance of renal perfusion. In these patients, administration of an NSAID may cause a dose-dependent reduction in prostaglandin formation and may precipitate overt renal decomposition. Patients at greatest risk of this reaction are those with impaired renal function, heart failure, liver dysfunction, nephrotic syndrome, those taking diuretics and the elderly. Such patients should be carefully monitored while receiving PONSTAN.

Discontinuation of NSAID therapy is typically followed by recovery to the pretreatment state. Since PONSTAN is eliminated primarily by the kidneys, the drug should not be administered to patients with significantly impaired renal functions (See section 4.3 - Contraindications).

Use in the Elderly

No data available.

Paediatric Use

Safety and effectiveness in children below the age of 14 years have not been established.

Effects on Laboratory Tests

A false positive reaction for urinary bile, using the diazo tablet test, may result after PONSTAN administration. If biliuria is suspected, other diagnostic procedures, such as the Harrison spot test, should be performed.

4.5 Interactions with other medicines and other forms of interactions

Aspirin

Mefenamic acid interferes with the anti-platelet effect of low-dose aspirin, and thus may interfere with aspirin's prophylactic treatment of CV disease.

Anticoagulants

The concurrent use of NSAIDs and warfarin has been associated with severe, sometimes fatal haemorrhage.

Mefenamic acid, like other non-steroidal anti-inflammatory agents, can inhibit platelet aggregation and may prolong prothrombin time in patients on warfarin therapy. Mefenamic acid has been shown to displace warfarin from protein binding sites and may enhance the response to oral anticoagulants.

NSAIDs, such as PONSAN, should be used in combination with warfarin, only if absolutely necessary. Concurrent administration of PONSAN with oral anticoagulant drugs requires frequent prothrombin time monitoring (See section 4.4 - Special warnings and precautions for use).

Anti-hypertensives

NSAIDs, such as mefenamic acid, can reduce the efficacy of anti-hypertensive drugs including diuretics, angiotensin-converting enzyme (ACE) inhibitors, angiotensin II antagonists (AIIA) and beta-blockers.

In patients with impaired renal function (e.g. dehydrated patients or elderly patients with compromised renal function), the co-administration of an ACE inhibitor or an AIIA and/or diuretics with a cyclo-oxygenase inhibitor can increase the deterioration of the renal function, including the possibility of acute renal failure, which is usually reversible. The occurrence of these interactions should be considered in patients taking mefenamic acid with an ACE inhibitor or an AIIA and/or diuretics.

Therefore, the concomitant administration of these drugs should be done with caution, especially in elderly patients. Patients should be adequately hydrated and the need to monitor renal function should be assessed before, and periodically during, concomitant treatment.

Corticosteroids

Concurrent use with NSAIDs may increase the risk of gastrointestinal ulceration or bleeding.

Cyclosporin or Tacrolimus

Concomitant administration with NSAIDs increases the risk of nephrotoxicity.

Hypoglycaemic Agents

There have been reports of changes in the effects of oral hypoglycaemic agents in the presence of NSAIDs. Therefore, mefenamic acid should be administered with caution in patients receiving insulin or oral hypoglycaemic agents.

Lithium

Mefenamic acid has produced an elevation of plasma lithium levels and a reduction in renal lithium clearance. Thus, when mefenamic acid and lithium are administered concurrently, patients should be observed carefully for signs of lithium toxicity.

Methotrexate

Caution is advised when methotrexate is administered concurrently with NSAIDs, including mefenamic acid, because NSAID administration may result in increased plasma levels of methotrexate, especially in patients receiving high doses of methotrexate.

4.6 Fertility, pregnancy and lactation

Effects on Fertility

Based on the mechanism of action, the use of NSAIDs may delay or prevent rupture of ovarian follicles, which has been associated with reversible infertility in some women. In

women who have difficulties conceiving or who are undergoing investigation of infertility, withdrawal of NSAIDs, including PONSTAN, should be considered.

Use in Pregnancy – Category C

Since there are no adequate and well-controlled studies in pregnant women, PONSTAN should be used only if the potential benefits to the mother justify the possible risks to the fetus. It is not known if mefenamic acid or its metabolites crosses the placenta.

NSAIDs given during the latter part of pregnancy may cause closure of the fetal ductus arteriosus, fetal renal impairment, inhibition of platelet aggregation, and delay labour and birth.

Because of the effects of drugs in this class (i.e. inhibitors of prostaglandin synthesis) on the fetal CV system (i.e. premature closure of the ductus arteriosus), the use of mefenamic acid in pregnant women is not recommended and should be avoided during the third trimester of pregnancy, including the last few days before expected birth.

Mefenamic acid inhibits prostaglandin synthesis which may result in prolongation of pregnancy and interference with labour when administered late in the pregnancy. Women on mefenamic acid therapy should consult their physician if they decide to become pregnant.

The inhibition of prostaglandin synthesis by NSAIDs may adversely affect pregnancy. Epidemiological studies suggest an increased risk of spontaneous abortion after use of prostaglandin synthesis inhibitors in early pregnancy. In animals, administration of prostaglandin synthesis inhibitors has been shown to result in increased pre- and post-implantation loss.

If used during second or third trimester of pregnancy, NSAIDs may cause fetal renal dysfunction which may result in reduction of amniotic fluid volume or oligohydramnios in severe cases. Such effects may occur shortly after treatment initiation. Pregnant women on mefenamic acid should be closely monitored for amniotic fluid volume.

Use in Lactation

Trace amounts of PONSTAN may be present in breast milk and transmitted to the nursing infant. Thus PONSTAN should not be taken by the nursing mother because of the effects of this class of drugs on the infant cardiovascular system.

4.7 Effects on ability to drive and use machines

The effects of this medicine on a person's ability to drive and use machines were not assessed as part of its registration. However, adverse effects of PONSTAN include dizziness, drowsiness and blurred vision which could affect the ability to drive or use machines (see Section 4.8 - Adverse effects (Undesirable effects)).

4.8 Adverse effects (undesirable effects)

Gastrointestinal Disorders

The most frequently reported adverse effects associated with the use of PONSTAN involve the gastrointestinal tract. In controlled studies for up to eight months, the following disturbances were reported in decreasing order of frequency; diarrhoea (approximately 5% of patients), nausea with or without vomiting, other gastrointestinal symptoms and abdominal pain. In certain patients, the diarrhoea was of sufficient severity to require discontinuation of the medication. Diarrhoea is usually dose related. It generally subsides on reduction of dosage and rapidly disappears on termination of therapy.

Other less frequently reported gastrointestinal effects were anorexia, cholestatic jaundice, colitis, enterocolitis, mild hepatic toxicity, hepatitis, hepatorenal syndrome, pyrosis, pancreatitis, steatorrhoea, flatulence, constipation, gastrointestinal inflammation, gastrointestinal ulceration (with and without haemorrhage) and gastrointestinal perforation.

Blood and Lymphatic System

Cases of auto-immune haemolytic anaemia have been associated with the continuous administration of PONSTAN for 12 months or longer. In such cases the Coombs test results are positive with evidence of both accelerated RBC production and RBC destruction. The process is reversible upon termination of PONSTAN administration.

Decreases in haematocrit have been noted in 2% to 5% of patients and primarily in those who have received prolonged therapy. Leukopenia, eosinophilia, thrombocytopenia purpura, agranulocytosis, pancytopenia, aplastic anaemia, bone marrow hypoplasia and platelet aggregation inhibition have also been reported.

Immune System

Anaphylaxis.

Metabolism and Nutrition

Glucose intolerance in diabetic patients, hyponatraemia and fluid retention.

Psychiatric Disorders

Nervousness.

Nervous System

Aseptic meningitis, convulsions, drowsiness, dizziness, headache, blurred vision and insomnia.

Eye and Ear Disorders

Eye irritation, reversible loss of colour vision and ear pain.

Cardiovascular System

Hypotension, hypertension and palpitations.

Respiratory and Thoracic System

Asthma and dyspnoea.

Skin and Subcutaneous Tissue Disorders

Angioedema, oedema of the larynx, erythema multiforme, perspiration, Lyell's syndrome (toxic epidermal necrolysis), Stevens-Johnson syndrome, dermatitis exfoliative, pruritus, urticaria, rash and facial oedema.

Renal and Urinary Disorders

As with other NSAIDs, renal failure including papillary necrosis has been reported. In elderly patients renal failure has occurred after taking PONENTAN for 2 to 6 weeks. The renal damage may not be completely reversible. Haematuria, dysuria, tubulointerstitial nephritis, glomerulonephritis and nephrotic syndrome have also been reported.

General Disorders and Administration Site Conditions

Oedema.

Investigations

Urobilinogen urine (false-positive) and liver function test abnormal.

Reporting Suspected Adverse Effects

Reporting suspected adverse reactions after registration of the medicinal product is important. It allows continued monitoring of the benefit-risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions at www.tga.gov.au/reporting-problems.

4.9 Overdose

Symptoms

Symptoms of overdose are related to the amount of drug ingested and range from gastrointestinal discomfort and diarrhoea to seizures, acute renal failure, confusional state, vertigo, hallucination, coma and death. Plasma levels of up to 210 µg/mL (therapeutic range 1 to 10 µg/mL) have been reported resulting in repeated generalised convulsions, but are not generally useful for evaluation and management of overdose.

Treatment

There is no specific antidote for mefenamic acid overdose. Treatment is symptomatic and supportive, including fluid replacement and IV access especially to patients who are dehydrated or unable to ingest adequate fluids. Avoiding intravascular fluid depletion will help prevent development of renal failure.

In cases of severe toxicity, activated charcoal may reduce absorption of the drug if given within one to two hours after ingestion. In patients who are not fully conscious or have impaired gag reflex, consideration should be given to administering activated charcoal via nasogastric tube ensuring that the airway is protected.

In clinically severe overdoses, full blood count, electrolytes, glucose, renal function, liver function tests, arterial blood gases and coagulation studies should be monitored for abnormalities.

Because mefenamic acid and its metabolites are firmly bound to plasma proteins, haemodialysis, haemoperfusion and peritoneal dialysis may be of little value.

For information on the management of overdose, contact the Poisons Information Centre on 13 11 26 (Australia).

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

PONSTAN has demonstrated analgesic, anti-inflammatory and antipyretic properties in human clinical studies and in classical animal test systems. These effects may be due to PONSTAN's dual action on prostaglandins. It inhibits the enzymes of prostaglandin synthetase and also antagonises the actions of prostaglandin at the receptor sites. These effects may also be responsible for its effectiveness in the treatment of primary dysmenorrhoea. The pain of primary dysmenorrhoea is thought to be due to increased abnormal uterine activity and uterine ischaemia, probably induced by release of $\text{PGF}_2\alpha$ or due to increase in the ratio of $\text{PGF}_2\alpha$: PGE_2 . Prostaglandins are also believed to be responsible, at least in some part, for the symptoms of menorrhagia.

Mechanism of Action

No data available.

Clinical Trials

No data available.

5.2 Pharmacokinetic properties

Absorption

Single and multiple studies have shown that PONSTAN usually reaches peak plasma levels 2 to 4 hours after oral administration with a half-life of 2 hours.

Distribution

Mefenamic acid and its metabolites are firmly bound to plasma proteins.

Metabolism

Mefenamic acid metabolism is predominantly mediated via cytochrome P450 CYP 2C9 in the liver. Patients who are known or suspected to be poor CYP2C9 metabolisers based on previous history/experience with other CYP2C9 substrates, should be administered mefenamic acid with caution as they may have abnormally high plasma levels due to reduced metabolic clearance.

Two distinct metabolic products, one a hydroxymethyl derivative and the other a carboxy derivative, have been identified in both plasma and urine. Mefenamic acid and its two metabolic derivatives become conjugated with glucuronic acid through an ester linkage which is alkali labile and are excreted principally in the urine, but also to some extent in the bile and faeces.

Excretion

Following a single dose, 67% of the total dose is excreted in the urine as unchanged drug or as one of the two metabolites. 20% to 25% of the dose is excreted in the faeces during the first three days.

5.3 Preclinical safety data

Genotoxicity

No data available.

Carcinogenicity

No data available.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

- Lactose monohydrate
- Titanium dioxide
- Gelatin
- Brilliant blue FCF
- Iron oxide yellow
- Carbon black

6.2 Incompatibilities

Incompatibilities were either not assessed or not identified as part of the registration of this medicine.

6.3 Shelf life

In Australia, information on the shelf life can be found on the public summary of the Australian Register of Therapeutic Goods (ARTG). The expiry date can be found on the packaging.

6.4 Special precautions for storage

Store below 30 °C.

6.5 Nature and contents of container

PONSTAN capsules are available in blister packs of 20 or 50 capsules and in bottles of 50 capsules. PONSTAN capsules are supplied in PVC/Al blister packs of 20 or 50 capsules.


6.6 Special precautions for disposal

In Australia, any unused medicine or waste material should be disposed of in accordance with local requirements.

6.7 Physicochemical properties

Mefenamic acid is one of the fenamate series of non-steroidal anti-inflammatory agents. It is a white or almost white, microcrystalline powder. The drug is almost insoluble in water, slightly soluble in ethanol (96 percent) and in methylene chloride. It dissolves in dilute solutions of alkali hydroxides.

Chemical Structure



Chemical name: N-(2,3, xylyl)-anthranilic acid

Molecular formula: C₁₅H₁₅NO₂

Molecular weight: 241.3

CAS Number

CAS Registry Number: 61-68-7.

7. MEDICINE SCHEDULE (POISONS STANDARD)

Pack of 20s: Schedule 2 (Pharmacy Medicine)

Pack of 50s: Schedule 4 (Prescription Only Medicine).

8. SPONSOR

Pfizer Australia Pty Ltd
Level 17, 151 Clarence Street
Sydney NSW 2000
Toll Free Number: 1800 675 229
www.pfizer.com.au

9. DATE OF FIRST APPROVAL

20 September 1991 (blister 20s and bottle 50s)

29 July 2014 (blister 50s)

10. DATE OF REVISION

20 December 2018

® Registered trademark.

Summary Table of Changes

Section changed	Summary of new information
All	Reformatting according to new Australian PI template
4.4	To update regarding the risk of GI complications in patients using concomitant antiplatelet drugs.