

MONOFEME(R)

Levonorgestrel and Ethinylloestradiol Tablets

Consumer Medicine Information

What is in this leaflet

This leaflet answers some common questions about Monofeme. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist.

All medicines have risks and benefits. Your doctor has weighed the risks of you taking Monofeme against the benefits this medicine is expected to have for you.

If you have any concerns about taking this medicine, ask your doctor or pharmacist.

Keep this leaflet with the medicine.

You may need to read it again.

What Monofeme is used for

Monofeme is an oral contraceptive. Monofeme tablets contain two hormones (levonorgestrel and ethinylloestradiol), which prevent you from becoming pregnant if taken correctly. They are similar to the hormones that your body normally produces.

Monofeme prevents pregnancy in several ways:

- * It inhibits the egg release by stopping it maturing
- * It changes the cervical mucus consistency making it difficult for the sperm to reach the egg
- * It changes the lining of the uterus making it less suitable for implantation.

Your doctor may have prescribed Monofeme for another reason. Ask your doctor if you have any questions about why Monofeme has been prescribed for you.

Monofeme is not habit-forming.

This medicine is available only with a doctor's prescription.

Before you take Monofeme

When you must not take Monofeme

Do not take Monofeme if:

- 1. You are allergic to any of the ingredients found in Monofeme**

The ingredients found in Monofeme are listed at the end of this leaflet.

Some of the symptoms of an allergic reaction may include rash, itching or hives on the skin, swelling of the face, lips, tongue or other parts of the body, shortness of breath, wheezing or troubled breathing.

- 2. You have or have had any medical conditions, especially the following:**

- * Blood clots in the legs (thrombophlebitis or deep vein thrombosis), lungs (pulmonary embolism) or eyes
- * Heart valve or heart rhythm disorders that may be associated with the formation of blood clots
- * Heart attack or stroke
- * Chest pain (angina pectoris)

- * High blood pressure which is uncontrolled
- * Some severe types of headache or migraine, including migraine with partial paralysis
- * Diabetes with blood vessel damage
- * Breast cancer or cancer of the lining of the womb, cervix or vagina, or you think you have these conditions
- * Unexplained vaginal bleeding
- * Liver tumour or liver disease
- * Inflammation of the pancreas, which is associated with very high blood levels of triglycerides (fatty substances).

If you are not certain whether these may apply to you, or you are worried by anything in this list, tell your doctor.

- 3. If you are pregnant or you think you are pregnant.**

Pregnancy must be excluded before you start taking Monofeme.

- 4. Monofeme is not for use in children or in postmenopausal women.**

- 5. The expiry date (EXP) printed on the pack has passed.**

Monofeme may have no effect at all, or worse, an entirely unexpected effect, if you take it after the expiry date.

- 6. The packaging is torn or shows signs of tampering.**

If this is the case, take the tablets back to your pharmacist.

Before you start to take Monofeme

You must have a thorough medical check-up, including a Pap smear, breast check, blood pressure check and urine test.

You must tell your doctor if:

- 1. You are allergic to any foods, dyes, preservatives or any other medicines.**
- 2. You smoke.**

Oral contraceptives increase your risk of having a stroke or heart attack. The risk of serious side effects on the heart and blood vessels is even greater if you smoke and take oral contraceptives. The risk increases with age and with heavy smoking (15 or more cigarettes per day), especially in women older than 35 years. If you take Monofeme, you should not smoke.

- 3. You have any other health problems, especially:**

- * Breast lumps, abnormal breast X-ray or mammogram
- * Diabetes
- * Heart disease
- * High cholesterol or blood fats
- * High blood pressure or you have a history of high blood pressure, including high blood pressure during pregnancy
- * Migraine or other headaches
- * Epilepsy
- * Depression
- * Gallbladder disease
- * Fluid retention or kidney disease
- * Asthma
- * Fibroids
- * Yellowing of the whites of the eyes or skin (jaundice) during pregnancy or during previous oral contraceptive use
- * Hereditary angioedema.

If you have any of these conditions you should have regular check-ups with your doctor to make sure that taking Monofeme is not making the conditions worse.

- 4. Anyone in your family has had blood clots in the deep veins of the legs, a stroke or heart attack or you have any of the following conditions:**

- * Obesity
- * Recent surgery or trauma
- * Recently had a baby
- * Lost a baby in the second trimester
- * Confined to bed rest for long periods

The risk of developing blood clots in the deep veins of your legs, which can break away and block a blood vessel elsewhere in your body, are increased if you have any of these conditions and use an oral contraceptive. Blood clots are a rare occurrence and can develop whether or not you are taking an oral contraceptive. They can also happen during pregnancy. The risk of having a blood clot is higher in oral contraceptive users than in non-users, but not as high as during pregnancy.

- 5. You plan to become pregnant or you think you are pregnant.**

Like most medicines, Monofeme is not recommended during pregnancy.

- 6. You are breast-feeding or plan to breast-feed.**

Like most medicines, Monofeme is not recommended while you are breast-feeding. Small amounts of oral contraceptives have been found in breast milk. It is not known what effect this may have on the baby. A decrease in milk supply may also occur.

If you have not told your doctor about any of the above, tell them before you start taking Monofeme.

Taking other medicines

Tell your doctor if you are taking any other medicines, including medicines you buy without a prescription from a pharmacy, supermarket or health food shop.

Some medicines may stop Monofeme from working properly. These

include medicines such as:

- * Rifampicin and rifabutin for the treatment of tuberculosis
- * Antibiotics such as ampicillin, other penicillins and tetracyclines
- * Anti-fungal agents such as griseofulvin
- * Barbiturates (phenobarbitone)
- * Medicines for epilepsy (such as phenytoin, primidone, carbamazepine and topiramate)
- * Ritonavir for the treatment of HIV infection
- * Modafinil used to treat excessive daytime sleepiness
- * St. John's wort, an ingredient in many medicines you can buy without a prescription from a pharmacy, health food shop or supermarket
- * Corticosteroids such as dexamethasone.

While you are taking any of these medicines and for the next 7 days after stopping them, you must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods). If you come to the end of the white tablets during these 7 days, start the next pack straight away. Skip the 7 red tablets.

If you take rifampicin and some other medicines, you may need to use additional non-hormonal contraception for four weeks after finishing the course of treatment.

Ask your doctor or pharmacist about how long you need to use additional non-hormonal contraception.

Some medicines may increase the levels of Monofeme in your blood, which may lead to unwanted side effects. These medicines include:

- * Atorvastatin used to treat high cholesterol
- * Indinavir for the treatment of HIV infection
- * Anti-fungal agents such as itraconazole and fluconazole
- * Paracetamol and ascorbic acid (Vitamin C).

Monofeme may also affect how well some other medicines work. These medicines include:

- * Cyclosporin used to prevent organ rejection
- * Theophyllines used for asthma and other breathing difficulties
- * Corticosteroids
- * Lamotrigine used for seizures.

If you have not told your doctor or pharmacist about any of the above, tell them before you start taking Monofeme.

If you are scheduled for any laboratory tests, tell your doctor you are taking Monofeme.

Some blood tests may be affected by taking Monofeme.

How to take Monofeme

Follow the directions on the blister pack.

If your doctor has prescribed Monofeme for some other purpose than contraception, follow their directions closely, even if they are not the same as on the pack.

When to take Monofeme

You must take Monofeme every day, even if you do not have sex very often.

Monofeme will work best if you do not miss any tablets and take it at the same time each day. Taking your tablet at the same time each day will also help you remember when to take the tablets.

It does not matter if you take Monofeme before or after food.

If you are concerned about this, please speak to your doctor or pharmacist.

How to take Monofeme

Swallow Monofeme with a glass of water.

Starting a hormonal contraceptive for the first time

To start taking Monofeme follow these steps:

1. On the first day of your menstrual bleed, take a white tablet that matches the day of the week from the grey shaded section of the blister pack.

You must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break.

2. Then take one white tablet each day, following the arrows so that you are taking the correct tablet for the day of the week until all 21 white tablets have gone.
3. Then take one red tablet each day for the next 7 days.
4. You will have a 'withdrawal' bleed, similar to having a period, during the week of red tablets.

If you do not bleed and there is any chance that you have not followed all the instructions in this leaflet, contact your doctor to check if you are pregnant.

Going on to further blister packs

1. On the day after your last red tablet, begin the next pack with a white tablet from the grey shaded section of the blister pack that matches the day of the week. Do this even if you are still bleeding.
2. Each new pack is started with a white tablet on the same day as the first pack, so that you have 21 days on white tablets, then 7 days on red tablets. There is no break between packs.
3. If you start the new pack later than the day after your last red tablet, you may have started a normal fertile cycle.

If you start late, you must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature

methods) until a white tablet has been taken daily for 7 days without a break.

Switching from a different combined oral contraceptive

Follow these steps if your current oral contraceptive contains an oestrogen and a progestogen:

1. Stop taking your current oral contraceptive after you have taken the last active tablet. If your current oral contraceptive pack also contains reminder tablets, do not take them.
2. The next day, take the first white Monofeme tablet from the grey shaded section that matches the day of the week.

You must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break.

3. Then take one white tablet each day following the direction of the arrows until all 21 white tablets have gone.
4. Then take one red tablet each day for the next 7 days.
5. You will have a 'withdrawal' bleed, similar to having a period, during the week of red tablets.

If you do not bleed and there is any chance that you have not followed all the advice in this leaflet, contact your doctor to check if you are pregnant.

Switching from a progestogen-only contraceptive

You can stop taking a progestogen-only contraceptive tablet any day and start taking Monofeme the next day, at the same time.

If you have been using a progestogen-only implant, start taking Monofeme on the day the implant is removed.

If you have been using a progestogen-only injection, start taking Monofeme on the day the next injection would be due.

In all cases start Monofeme by taking a white tablet from the grey shaded section that matches the day of the week.

You must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break.

After having a baby

If you have just had a baby, talk to your doctor before you start taking Monofeme.

After a miscarriage or abortion

Your doctor will advise you how to take Monofeme after a miscarriage or abortion.

How long to take Monofeme

Your doctor may prescribe Monofeme for long periods, until you no longer need or want contraception.

If you are not sure how long you should be taking Monofeme, ask your doctor.

If you forget to take your tablets

If you forget to take Monofeme every day it may not work as well in protecting you from becoming pregnant.

Do not try to make up for missed doses by taking more than one tablet at a time.

FORGETTING ONE WHITE TABLET

- 1. If you forget one white tablet but it is less than 12 hours late, take the missed tablet immediately. Take the next tablet at your usual time, even**

if this means taking two white tablets in one day.

If you do not take the missed tablet within 12 hours, Monofeme may not work as well in protecting you from becoming pregnant.

- 2. If one white tablet is missed and is more than 12 hours, take the last tablet as soon as you remember and the next white tablet at the usual time.**

Continue to take tablets at your usual time but you must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break. If you come to the end of the white tablets during the 7 days after a missed tablet, start the next pack straight away. Skip the 7 red tablets.

FORGETTING TWO WHITE TABLETS

- 1. If you forget two white tablets, take the last missed tablet as soon as you remember and the next tablet at the usual time. Continue to take tablets at your usual time but you must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break. If you come to the end of the white tablets during the 7 days after the missed tablets, start the next pack straight away. Skip the 7 red tablets.**

You may not have a withdrawal bleed until the end of this new pack, but you may have spotting or breakthrough bleeding when taking the white tablets. If you don't have a withdrawal bleed at the end of the new pack, contact your doctor, who

must make sure you are not pregnant before you start the next pack.

FORGETTING THREE OR MORE WHITE TABLETS

Contact your doctor for advice on what to do.

FORGETTING A RED TABLET

- 1. If you miss one or more red tablets, leave them in the pack and do not worry.**
- 2. However, if you miss red tablets and then forget to start the next pack on time, start as soon as you remember by taking a white tablet that matches the day of the week from the grey shaded section. You must also use an additional non-hormonal method of contraception (such as condoms or a diaphragm but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break.**

If you are not sure what to do, ask your doctor or pharmacist.

If you are having trouble remembering to take Monofeme, ask your pharmacist for some hints.

If you vomit or have diarrhoea after taking Monofeme

If you have vomiting or diarrhoea within 3 to 4 hours after taking a white tablet, you must use an additional non-hormonal method of contraception (such as condoms or a diaphragm, but not the rhythm or temperature methods) until a white tablet has been taken daily for 7 days without a break. If you come to the end of the white tablets during these 7 days, start the next pack straight away. Skip the 7 red tablets.

The tablet may not have time to be absorbed properly and may not protect you from becoming pregnant.

If you have vomiting or diarrhoea after taking a red tablet, do not worry.

If you take too much (overdose)

Immediately telephone your doctor or the Poisons Information Centre (Australia telephone 13 11 26, New Zealand telephone 0800 764 766) for advice, or go to Accident and Emergency at your nearest hospital if you think you or anyone else may have taken too much Monofeme. Do this even if there are no signs of discomfort or poisoning. You may need urgent medical attention.

If you take too much Monofeme, some of the symptoms you may have include:

- * Feeling sick or vomiting
- * Dizziness
- * Feeling sleepy or tired.

Women may also experience menstrual bleeding.

While you are taking Monofeme

Things you must do

Tell all doctors, dentists and pharmacists who are treating you that you are taking Monofeme.

If you are about to start taking any new medicines, tell the doctor or pharmacist that you are taking Monofeme.

If you become pregnant while taking Monofeme, see your doctor immediately.

If you miss a period and you have taken your tablets correctly, continue taking your tablets as you would normally.

Sometimes you might not have a menstrual period while taking Monofeme.

If you miss a period and you have not taken your tablets correctly, keep taking your tablets and see your doctor immediately.

Not taking your tablets correctly includes missing one or more tablets

or starting a new pack later than you should have.

If you miss two menstrual periods, stop taking your tablets and see your doctor, even if you have taken the tablets correctly. You must use a non-hormonal method of contraception, (such as condoms or a diaphragm) during this time.

Your doctor should make sure you are not pregnant before you start taking Monofeme again.

Have regular check ups from your doctor, including a Pap smear.

Oral contraceptives should not be prescribed for longer than one year without your doctor carrying out a check-up. Your doctor will advise you how often you need a Pap smear. A Pap smear can detect abnormal cells lining the cervix. Sometimes abnormal cells can progress to cervical cancer. The most important risk factor for cervical cancer is persistent human papillomavirus (HPV) infection. However, cervical cancer has been reported to occur more often in women using an oral contraceptive for a long time. This finding may not be caused by the oral contraceptive, but may be related to sexual behaviour and other reasons.

Perform regular breast self-examination.

Risk factors for the development of breast cancer include increasing age, family history, obesity, never having had a baby, and late age for first full-term pregnancy. Breast cancer has also been found slightly more often in women who use oral contraceptives than in women of the same age who do not use them. This slight increase in the number of breast cancer cases gradually disappears during the course of the 10 years after stopping use of oral contraceptives. It is not known whether the oral contraceptive causes the difference. It may be that the women were examined more often, so that the breast cancer was noticed earlier.

If you are concerned about contracting a sexually transmitted disease (STD), ask your partner to wear a condom when having sexual intercourse with you.

Monofeme will not protect you from HIV-AIDS or any other sexually transmitted diseases such as chlamydia, genital herpes, genital warts, gonorrhoea, hepatitis B, human papilloma virus and syphilis. To help protect yourself from STDs, you need to use a barrier contraceptive such as a condom.

Tell your doctor you are using Monofeme at least 4 weeks before any planned hospitalisation or surgery.

Your doctor may tell you to stop taking Monofeme several weeks before surgery or at the time of immobilisation. Your doctor will tell you when you can start taking Monofeme after you are back on your feet.

To avoid pregnancy during this time you must use a non-hormonal method of contraception such as condoms or a diaphragm.

Things you must not do

Do not give Monofeme to anyone else even if they have the same condition as you.

Do not use Monofeme to treat any other complaints unless your doctor tells you to.

Do not stop taking Monofeme, or change the dosage, without checking with your doctor.

If you stop taking Monofeme or do not take a tablet every day, without using another form of contraception, you may become pregnant.

Side Effects

Tell your doctor or pharmacist as soon as possible if you do not feel well while taking Monofeme.

When used correctly, Monofeme is

an effective contraceptive, but may have unwanted side effects in some people. All medicines can have side effects. Sometimes they are serious, most of the time they are not.

Ask your doctor or pharmacist to answer any questions you have.

Tell your doctor immediately, or go to accident and emergency at your nearest hospital if you notice any of the following:

- * Sharp chest pain, coughing of blood, or sudden shortness of breath
- * Pain in the calf muscle area
- * Crushing chest pain or heaviness in the chest
- * Sudden severe headache or vomiting, dizziness or fainting, disturbances of vision or speech, weakness, or numbness in an arm or leg
- * Sudden changes or loss of vision
- * Breast lumps
- * Severe pain or tenderness in the stomach area
- * Jaundice or a yellowing of the skin or eyeballs, often with fever, fatigue, loss of appetite, dark coloured urine or light coloured bowel movements. Taking oral contraceptives has been associated with an increased risk of having a benign liver tumour and, in very rare cases, liver cancer. The risk appears to increase the longer oral contraceptives are taken.
- * Migraine headaches for the first time
- * More frequent migraines if you already suffer from them
- * Itchy rash
- * You are an epileptic and your fits become more frequent
- * Rise in blood pressure
- * Swelling around eyes or mouth
- * Bloody diarrhoea, abdominal pain or tenderness, fever, nausea or vomiting.

Whilst these side effects are rare, they are serious. You may need urgent medical attention or

hospitalisation.

Tell your doctor if you notice any of the following.

Do not be alarmed by this list of possible side effects. You may not experience any of them.

Reproductive or breast problems such as:

- * Changes in bleeding patterns, including breakthrough bleeding/spotting
- * Painful periods
- * Missed periods, but if you have not taken Monofeme as directed you should check whether you are pregnant
- * Changes in mucus from the vagina
- * Changes in the cervix
- * Vaginal thrush (candida)
- * Breast pain, tenderness, enlargement, possible milk secretion
- * Changes in sex drive.

Stomach problems such as:

- * Nausea or vomiting
- * Abdominal pain, cramps or bloating.

Difficulties thinking or working because of:

- * Mood changes, including depression
- * Headache, including migraines
- * Nervousness
- * Dizziness
- * Contact lenses becoming uncomfortable to wear.

Changes to your appearance such as:

- * Weight change (increase or decrease) or changes in appetite
- * Swelling of the hands, ankles or feet
- * Acne
- * Rash
- * Darkening of the skin, which may persist after stopping Monofeme
- * Loss of scalp hair
- * Increase in body hair.

Worsening of conditions you may already have such as:

- * Chorea
- * Porphyria
- * Systemic lupus erythematosus
- * Varicose veins
- * Gallbladder disease
- * Hereditary angioedema.

Other side effects not listed above may also occur in some patients. Tell your doctor if you notice anything else that is making you feel unwell, even if it is not on this list.

After stopping Monofeme

If your periods do not return within 2 to 3 months of stopping Monofeme tell your doctor.

Some women have short-term problems getting pregnant after stopping Monofeme, especially if they had irregular menstrual cycles before starting to use an oral contraceptive.

If you are planning to become pregnant after stopping Monofeme, use a non-hormonal method of contraception such as condoms or a diaphragm for 3 months before trying to get pregnant.

Ask your doctor or pharmacist for advice about taking folate if you plan to become pregnant.

After taking Monofeme

Storage

Keep your tablets in the blister pack until it is time to take them.

If you take the tablets out of the blister pack they may not keep well.

Keep Monofeme in a cool, dry place where the temperature stays below 25 degrees C and is away from light.

Do not store Monofeme or any other medicine, in a bathroom or near a sink.

Do not leave Monofeme in the car on hot days or on window sills.

Heat and dampness can destroy some medicines.

Keep Monofeme where children cannot reach it.

A locked cupboard at least one-and-a-half metres above the ground is a good place to store medicines.

Disposal

If your doctor tells you to stop taking Monofeme, or the tablets have passed their expiry date, ask your pharmacist what to do with any left over.

Product description

What it looks like

In Australia Monofeme comes in a 4-week box containing 1 blister pack and a 16-week box containing 4 blister packs. In New Zealand Monofeme comes in a 4-week box containing 1 blister pack and a 12-week box containing 3 blister packs. Each blister pack contains 21 white hormone tablets and 7 red non-hormonal tablets. The blister pack is marked with days of the week next to each tablet.

Ingredients

Each white tablet contains 150 micrograms of levonorgestrel and 30 micrograms of ethinyloestradiol as the active ingredients.

Each red tablet contains no active ingredients.

The white and red tablets also contain the following inactive ingredients:

- * Lactose
- * Maize starch
- * Povidone
- * Talc
- * Magnesium stearate

- * Sucrose
- * Macrogol 6000
- * Calcium carbonate
- * Glycol montanate (red tablets only)
- * White beeswax (white tablets only)
- * Carnauba Wax (white tablets only)

The red tablets also contain the colouring agents:

- * Brilliant scarlet 4R
- * Erythrosine

Monofeme does not contain gluten, tartrazine or any other azo dyes.

Supplier

Australia

Monofeme is supplied by:

Wyeth Australia Pty Limited,

ABN 16 000 296 211

17-19 Solent Circuit,

Norwest Business Park,

Baulkham Hills NSW 2153

For further information please contact Wyeth on toll-free 1800 500 498 or E-mail: medinfo@wyeth.com

New Zealand

For all customer enquiries please contact Wyeth on toll-free 0800 447 400 or Email: medinfo@wyeth.com

Distributed in New Zealand by:

Pharmacy Retailing (NZ) Limited

58 Richard Pearce Drive, Airport Oaks, Auckland

Australian Registration Number:

AUST R 153764

This leaflet was prepared on 10 November 2009.

(R) Registered Trade Mark